ELSEVIER

www.elsevier.nl/locate/jorganchem

Journal of Organometallic Chemistry 621 (2001) 359

Journal of Organometallic Chemistry

Author Index of Volume 621

Alt, H.G., 304
Amor, F., 3
Angelici, R.J., 55
Apperley, D.C., 254
Auner, N., 10

Back, S., 143
Barbaro, P., 26
Barz, M., 153
Baumann, W., 89
Bäumer, U., 261
Beck, W., 109
Bellassoued, M., 284
Beller, M., 70
Bergman, R.G., 242
Bianchini, C., 26
Blaser, H.-U., 34
Bosl, G., 352
Braunstein, P., 218
Buser, H.-P., 34

Cadierno, V., 39
Calléja, G., 46
Caporusso, A.M., 246
Cerveau, G., 46
Chelain, E., 284
Chen, J., 55
Chisholm, M.H., 66
Comte, V., 284
Corriu, R.J.P., 46

Davies, N.A., 254
Döbler, C., 70
Dorfer, A., 173
Dötz, K.H., 77
Dreier, T., 197
Drexler, H.-J., 89

Ederer, T., 109
Eibl, S., 166
Elbertse, G., 337
Erker, G., 197

Faure, M., 103
Fehn, A., 109
Fehring, V., 120
Fischer, C., 89
Fischer, H., 344
Fischer, R.D., 254
Fokken, S., 3
Franció, G., 130
Fröhlich, R., 197
Frosch, W., 143

Gamasa, M.P., 39
Garrier, E., 284
Gimeno, J., 39
Girgsdies, F., 317
Glas, H., 153
Gonçalves, I.S., 207
Grasmann, M., 10
Grassert, I., 158
Grosche, M., 177, 184

Haage, K., 120
Harris, R.K., 254
Häusel, R., 34
Heller, D., 89
Herberhold, M., 166
Herrmann, W.A., 1, 177, 184
Holz, J., 120

Jäger, K., 173
Jaitner, P., 173
Jalett, H.-P., 34

Kadyrov, R., 120
Kaul, F.A.R., 177, 184
Kempe, R., 261
Kleij, A.W., 190
Klein Gebbink, R.J.M., 190
Kleinhenn, T., 3
Kühn, F.E., 207

Lang, H., 143
Leitner, W., 130
Lopes, A.D., 207
Ludwig, M., 120
Lutz, M., 190
Lynn, M.A., 66

Maiwald, S., 327
Mashima, K., 224
Mattheis, C., 218
Mehltretter, G.M., 70
Mickoleit, M., 261
Mihalios, D., 177, 184
Mihan, S., 109
Milius, W., 166

Nakamura, A., 224
Nieger, M., 77
Noltemeyer, M., 310
Nöth, H., 231

Oehme, G., 158
Oehme, H., 261
Okuda, J., 3
Otto, F., 77

Paisner, S.N., 242
Papadogianakis, G., 337
Pertici, P., 246
Pillinger, M., 207
Pitzalis, E., 246
Poll, E.-M., 254
Pötter, M., 261
Puchta, G.T., 177, 184

Razavi, A., 267
Robl, C., 352
Rodriguez-Borges, J.E., 207
Roesky, H.W., 310
Roesky, P.W., 277
Romão, C.C., 207
Royo, P., 299
Rudler, H., 284

Sánchez-Nieves, J., 299
Santos, A.M., 207
Scapacci, G., 26
Schlegel, A., 231
Schmidt, R., 304
Schneider, H., 177, 184
Schormann, M., 310
Schumann, H., 317
Schütze, J.-U., 254
Schwarzhans, K.-E., 173
Selke, R., 120
Sheldon, R.A., 337
Sieler, J., 327
Spaniol, T.P., 3
Spannenberg, A., 89
Spek, A.L., 190
Spindler, F., 34
Stoeckli-Evans, H., 103
Sundermeier, U., 70
Süss-Fink, G., 103
Suter, M., 231

Taube, R., 327
Tesouro Vallina, A., 103
Thewalt, U., 267
Thiel, W.R., 153

Vaissermann, J., 284
van Koten, G., 190
Varkey, S.P., 310
Verrazzani, A., 246
Verspui, G., 337
Vitulli, G., 246

Wassermann, B.C., 317
Weißenbach, K., 344
Weigand, W., 352
Wernik, S., 317
Wittmann, K., 130
Wünsch, R., 352

PII: S0022-328X(01)00691-X

ELSEVIER

www.elsevier.nl/locate/jorganchem

Journal of Organometallic Chemistry 621 (2001) 361–367

Journal of Organometallic Chemistry

Subject Index of Volume 621

Air
Dihydroxylation of olefins using air as the terminal oxidant, 70
Alkeneinylhalogensilanes lithium-*tert*-butyl
Silaheterocyclen XXXVII. Die Kombination Alkeninylhalogensilan/Lithium-*tert*-Butyl als Synthesebaustein zum Aufbau von Silaheterocyclen mit kumulierten CC-Doppelbindungen, 10
Alkoxides
Preparation of bistrimethylsilylmethylniobiumtetrafluoride and the application of KHF_2 and n-Bu_4NHF_2 as fluorinating reagents, 310
Alkylation
Ligands with cycloalkane backbones VIII. Steric effects in the ligand accelerated enantioselective alkylation of benzaldehyde, 153
Alkyl polyglycosides
Synthesis of achiral, but unsymmetric, seven-membered rhodium(I)-chelates for hydrogenation in the chiral environment of alkyl polyglucoside micelles, 120
Alkyne cyclotrimerisation
Stoichiometric cyclotrimerisation of chiral alkynes at a ruthenium centre: preparation of optically active (η^6-arene)(η^4-cycloocta-1,5-diene)ruthenium(0) complexes, 246
Alkynyl complexes
Stereoselective synthesis of chiral terminal (*E*)-1,3-enynes derived from the aldehydes (1*R*)-(−)-myrtenal and (*S*)-(−)-perillaldehyde using the alkynyl–phosphonio complex [Ru{C≡CCH$_2$(PPh$_3$)}(η^5-C$_9$H$_7$)(PPh$_3$)$_2$][PF$_6$] as synthon, 39
Allenylidene complexes
Stereoselective synthesis of chiral terminal (*E*)-1,3-enynes derived from the aldehydes (1*R*)-(−)-myrtenal and (*S*)-(−)-perillaldehyde using the alkynyl–phosphonio complex [Ru{C=CCH$_2$(PPh$_3$)}(η^5-C$_9$H$_7$)(PPh$_3$)$_2$][PF$_6$] as synthon, 39
Allyl
Komplexkatalyse: LVII. Vereinfachte Synthese des Nd(π-C$_3$H$_5$)$_3$·C$_4$H$_8$O$_2$ nach der Grignard-Methode und Darstellung der neuen Allylneodym(III)-Komplexe [Nd(π-C$_3$Me$_5$)(π-C$_3$H$_5$)$_2$·C$_4$H$_8$O$_2$] und [Nd(π-C$_3$H$_5$)Cl(THF)$_3$]B(C$_6$H$_5$)$_4$·THF als Präkatalysatoren für die stereospezifische Butadienpolymerisation, 327
Aluminium
Synthese und Charakterisierung chiraler Organogallium- und -aluminiumalkoholate mit intramolekularer Stickstoffstabilisierung. Röntgenstrukturanalyse von Dimethylgallium-(*S*)-2-dimethylamino-3-methyl-1-butanolat, Dimethylgallium-(*S*)-2-dimethylamino-3-phenyl-1-propanolat und Dimethylgallium-(*R*)-2-dimethylamino-1-butanolat, 317
Amine alcoholate
Synthese und Charakterisierung chiraler Organogallium- und -aluminiumalkoholate mit intramolekularer Stickstoffstabilisierung. Röntgenstrukturanalyse von Dimethylgallium-(*S*)-2-dimethylamino-3-methyl-1-butanolat, Dimethylgallium-(*S*)-2-dimethylamino-3-phenyl-1-propanolat und Dimethylgallium-(*R*)-2-dimethylamino-1-butanolat, 317
α-Amino acids
Metallkomplexe mit biologisch wichtigen Liganden, CXXVII: Halbsandwich-Komplexe von Ruthenium, Rhodium und Iridium sowie phosphanhaltige Palladium- und Platin(II)-Komplexe mit Schiff-Basen aus α-Aminosäureanionen und Aldehyden, 109
Aminotroponiminate
Cyclooctatetraene complexes of yttrium and the lanthanides with chiral and achiral aminotroponiminates, 277
Amphiphiles
Aqueous micellar and non-micellar effects during the asymmetric hydrogenation of dehydroamino acid derivatives: influence of amphiphiles on enantioselectivity and α-CH/CD exchange, 158
Aqueous biphase catalysis
Transition metal complexes with the C_1-symmetric diphosphines (*R*)-(*R*)-3-benzyl-2,4-bis(diphenylphosphino)pentane and (*R*)-(*R*)-3-benzyl(*p*-sulphonate)-2,4-bis(diphenylphosphino)-pentane sodium salt. Applications to enantioselective catalysis in different phase systems, 26
Arene complexes
Stoichiometric cyclotrimerisation of chiral alkynes at a ruthenium centre: preparation of optically active (η^6-arene)(η^4-cycloocta-1,5-diene)ruthenium(0) complexes, 246
Arene ligand
Catalytic hydrogenation of aromatics under biphasic conditions: isolation and structural characterisation of the cluster intermediate [(η^6-C$_6$Me$_6$)$_2$(η^6-C$_6$H$_6$)Ru$_4$(μ_2-H)$_2$(μ_2-OH)(μ_3-O)]$^+$, 103
Aromatic hydrogenation
Catalytic hydrogenation of aromatics under biphasic conditions: isolation and structural characterisation of the cluster intermediate [(η^6-C$_6$Me$_6$)$_2$(η^6-C$_6$H$_6$)Ru$_4$(μ_2-H)$_2$(μ_2-OH)(μ_3-O)]$^+$, 103
Aryl ethers
Meta, *para* and *meta*, *ortho* double *exo* nucleophilic additions of trimethylsilylester enolates derived from saturated and unsaturated carboxylic acids to tricarbonylchromium complexes of aryl ethers: dearomatizing cyclization to lactones, 284
Asymmetric catalysis
Transition metal complexes with the C_1-symmetric diphosphines (*R*)-(*R*)-3-benzyl-2,4-bis(diphenylphosphino)pentane and (*R*)-(*R*)-3-benzyl(*p*-sulphonate)-2,4-bis(diphenylphosphino)-pentane sodium salt. Applications to enantioselective catalysis in different phase systems, 26
Asymmetric dihydroxylation
Dihydroxylation of olefins using air as the terminal oxidant, 70
Asymmetric hydrogenation
Aqueous micellar and non-micellar effects during the asymmetric hydrogenation of dehydroamino acid derivatives: influence of amphiphiles on enantioselectivity and α-CH/CD exchange, 158

PII: S0022-328X(01)00692-1

Synthesis of achiral, but unsymmetric, seven-membered rhodium(I)-chelates for hydrogenation in the chiral environment of alkyl polyglucoside micelles, 120

Binaphtholato aluminates
A study of Noyori's reagent, 231
Bis(alkynyl) titanocenes
Alkinstabilisierte monomere Organo-Kupfer(I)-Verbindungen als Organyl-Transfer Reagenzien, 143
Bisiminopyridines
Trimethylammonium[tetrakis(pentafluorophenoxy)borate], a safe way to synthesize novel co-catalysts for olefin polymerization, 184
Bis[2,2,6,6-tetramethylpiperidyl-(1)]-disulfide
Improved synthesis of the radical source bis[2,2,6,6-tetramethylpiperidyl-(1)]-disulfide and photoinduced reactions with $M_2(CO)_{10}$ (M = Mn, Re), 173
Bonding
The electronic structure and bonding in $W_2(\mu\text{-}H)_2(O^iPr)_4(DMPE)_2$ and a comparison with $Mo_2(O^iPr)_4(DMPE)_2$, 66
Borates
Syntheses and properties of the novel co-catalysts *N,N*-dimethylanilinium- and trityl{tetrakis[4-(trifluoromethyl)-2,3,5,6-tetrafluorophenyl]borate}, 177
Trimethylammonium[tetrakis(pentafluorophenoxy)borate], a safe way to synthesize novel co-catalysts for olefin polymerization, 184
Butadiene
Komplexkatalyse: LVII. Vereinfachte Synthese des $Nd(\pi\text{-}C_3H_5)_3\text{-}C_4H_8O_2$ nach der Grignard-Methode und Darstellung der neuen Allylneodym(III)-Komplexe $[Nd(\pi\text{-}C_3Me_5)(\pi\text{-}C_3H_5)_2\text{-}C_4H_8O_2]$ und $[Nd(\pi\text{-}C_3H_5)Cl(THF)_3]B(C_6H_5)_4\cdot THF$ als Präkatalysatoren für die stereospezifische Butadienpolymerisation, 327

Carbamate compounds
Insertion of carbon dioxide and isocyanide into tantalum–amide and tantalum–methyl bonds, 299
Carbene complexes
Synthesis of chiral β-aminoalcohol-substituted carbene complexes of manganese and influence of the chiral carbene ligand on the diastereoselectivity of the CO/PR_3 exchange, 344
Carbon dioxide
Insertion of carbon dioxide and isocyanide into tantalum–amide and tantalum–methyl bonds, 299
Carbosilanes
Synthesis and characterization of platinum(II)-terminated dendritic carbosilanes: X-ray crystal structure of the model species $[PtCl(C_6H_3\{CH_2NMe_2\}\text{-}2\text{-}SiMe_2\text{-}5)(PPh_3)]$, 190
Catalysis
Unbridged metallocene dichloride complexes with mono-substituted indenyl ligands and their application for the polymerization of propene, 304
Catalyst
Komplexkatalyse: LVII. Vereinfachte Synthese des $Nd(\pi\text{-}C_3H_5)_3\text{-}C_4H_8O_2$ nach der Grignard-Methode und Darstellung der neuen Allylneodym(III)-Komplexe $[Nd(\pi\text{-}C_3Me_5)(\pi\text{-}C_3H_5)_2\text{-}C_4H_8O_2]$ und $[Nd(\pi\text{-}C_3H_5)Cl(THF)_3]B(C_6H_5)_4\cdot THF$ als Präkatalysatoren für die stereospezifische Butadienpolymerisation, 327
CC-double bonds
Silaheterocyclen XXXVII. Die Kombination Alkeninylhalogensilan/Lithium-*tert*-Butyl als Synthesebaustein zum Aufbau von Silaheterocyclen mit kumulierten CC-Doppelbindungen, 10

C–H activation
Synthesis and characterization of platinum(II)-terminated dendritic carbosilanes: X-ray crystal structure of the model species $[PtCl(C_6H_3\{CH_2NMe_2\}\text{-}2\text{-}SiMe_2\text{-}5)(PPh_3)]$, 190
Chelate ligands
Ligands with cycloalkane backbones VIII. Steric effects in the ligand accelerated enantioselective alkylation of benzaldehyde, 153
Chiral complexes
Synthesis of chiral β-aminoalcohol-substituted carbene complexes of manganese and influence of the chiral carbene ligand on the diastereoselectivity of the CO/PR_3 exchange, 344
Chiral ligands
Chelating behaviour of a phosphinoalcohol leading to a stable alcohol palladium complex, 218
Chiral bis(oxazoline) and pyridyl alcoholate dioxo-molybdenum(VI) complexes: synthesis, characterization and catalytic examinations, 207
Cluster
Catalytic hydrogenation of aromatics under biphasic conditions: isolation and structural characterisation of the cluster intermediate $[(\eta^6\text{-}C_6Me_6)_2(\eta^6\text{-}C_6H_6)Ru_3(\mu_3\text{-}H)_2(\mu_2\text{-}OH)(\mu_3\text{-}O)]^+$, 103
Co-catalyst
Syntheses and properties of the novel co-catalysts *N,N*-dimethylanilinium- and trityl{tetrakis[4-(trifluoromethyl)-2,3,5,6-tetrafluorophenyl]borate}, 177
Co-catalysts
Trimethylammonium[tetrakis(pentafluorophenoxy)borate], a safe way to synthesize novel co-catalysts for olefin polymerization, 184
Copper (I) cyanide
Self-assembly of $[Cu(CN)_4]^{3-}$ ions with cationic $\{Me_3Sn\}^+$ or $\{Me_2Sn(CH_2)_3SnMe_2\}^{2+}$ fragments in the presence of a $^nBu_4N^+$ template, 254
Counter-ions
Synthesis of achiral, but unsymmetric, seven-membered rhodium(I)-chelates for hydrogenation in the chiral environment of alkyl polyglucoside micelles, 120
Crystal structure
Cationic derivatives of $RhCl[P(\eta^2\text{-}C_7H_{11})_3]$. An intramolecular Diels–Alder rearrangement in the tripodal ligand tri(1-cyclohepta-2,4,6-trienyl)phosphane, $P(C_7H_9)_3$, 166
Crystal structures
Preparation and crystal structures of the complexes $(\eta^5\text{-}C_5H_3TMS\text{-}CMe_2\text{-}\eta^5\text{-}C_{13}H_8)MCl_2$ and $[3,6\text{-}di^tBuC_{13}H_6\text{-}SiMe_2\text{-}N^tBu]MCl_2$ (M = Hf, Zr or Ti): mechanistic aspects of the catalytic formation of a isotactic–syndiotactic stereoblock-type polypropylene, 267
C_1-symmetric diphosphines
Transition metal complexes with the C_1-symmetric diphosphines (*R*)-(*R*)-3-benzyl-2,4-bis(diphenylphosphino)pentane and (*R*)-(*R*)-3-benzyl(*p*-sulphonate)-2,4-bis(diphenylphosphino)-pentane sodium salt. Applications to enantioselective catalysis in different phase systems, 26
1,5-Cyclooctadiene complexes
Stoichiometric cyclotrimerisation of chiral alkynes at a ruthenium centre: preparation of optically active $(\eta^6\text{-arene})(\eta^4\text{-cycloocta-1,5-diene})ruthenium(0)$ complexes, 246
Cyclooctatetraene
Cyclooctatetraene complexes of yttrium and the lanthanides with chiral and achiral aminotroponiminates, 277
Cyclopentadienyl ligand
Mono(cyclopentadienyl)titanium complexes containing a sulfide-bridged bis(phenolato) ligand. Molecular structure of $Ti\{2,2'\text{-}S(OC_6H_2\text{-}4\text{-}Me\text{-}6\text{-}^tBu)_2\}(\eta^5\text{-}C_5H_5)Cl$, 3

Dearomatization
Meta, para and *meta, ortho* double *exo* nucleophilic additions of trimethylsilylester enolates derived from saturated and unsaturated carboxylic acids to tricarbonylchromium complexes of aryl ethers: dearomatizing cyclization to lactones, 284
Dendrimers
Synthesis and characterization of platinum(II)-terminated dendritic carbosilanes: X-ray crystal structure of the model species $[PtCl(C_6H_3\{CH_2NMe_2\}\text{-}2\text{-}SiMe_3\text{-}5)(PPh_3)]$, 190
Deuteration
Aqueous micellar and non-micellar effects during the asymmetric hydrogenation of dehydroamino acid derivatives: influence of amphiphiles on enantioselectivity and α-CH/CD exchange, 158
Diastereoselectivity
Diastereoselective oxidative addition of cyclic thiosulfinates to platinum(0) compounds: chiral platinum(II) complexes with sulfenato ligands Crystal structures of $S(O)–CH_2–CH(OAc)–CH(OAc)–CH_2–S$ and dppePt-$[S(O)–(CH_2)_4–S]$: Part 16. Metal complexes of functionalized sulfur-containing ligands, 352
Synthesis of chiral β-aminoalcohol-substituted carbene complexes of manganese and influence of the chiral carbene ligand on the diastereoselectivity of the CO/PR_3 exchange, 344
Diels–Alder cycloaddition2
Cationic derivatives of $RhCl[P(\eta^2\text{-}C_2H_3)_3]_2$. An intramolecular Diels–Alder rearrangement in the tripodal ligand tri(1-cyclohepta-2,4,6-trienyl)phosphane, $P(C_7H_7)_3$, 166
Diene complexes
'*Supine*' or '*prone*' ligands: geometric preference of conjugated diene, 1-azadiene, and 1,4-diazadiene ligands on half-metallocene complexes of early transition metals, 224
Dienol
Meta, para and *meta, ortho* double *exo* nucleophilic additions of trimethylsilylester enolates derived from saturated and unsaturated carboxylic acids to tricarbonylchromium complexes of aryl ethers: dearomatizing cyclization to lactones, 284
Diethylzinc
Ligands with cycloalkane backbones VIII. Steric effects in the ligand accelerated enantioselective alkylation of benzaldehyde, 153
Ditopic ligands
Chelating behaviour of a phosphinoalcohol leading to a stable alcohol palladium complex, 218

Electronic structures
The electronic structure and bonding in $W_2(\mu\text{-}H)_2(O^iPr)_4(DMPE)_2$ and a comparison with $Mo_2(O^iPr)_4(DMPE)_2$, 66
Enantioselective catalysis
Ligands with cycloalkane backbones VIII. Steric effects in the ligand accelerated enantioselective alkylation of benzaldehyde, 153
Enantioselective hydroformylation
Highly efficient enantioselective catalysis in supercritical carbon dioxide using the perfluoroalkyl-substituted ligand (*R,S*)-3-H^2F^6-BINAPHOS, 130
Enantioselective hydrogenation
Highly efficient enantioselective catalysis in supercritical carbon dioxide using the perfluoroalkyl-substituted ligand (*R,S*)-3-H^2F^6-BINAPHOS, 130
Tunable ferrocenyl diphosphine ligands for the Ir-catalyzed enantioselective hydrogenation of *N*-aryl imines, 34
1,3-Enynes
Stereoselective synthesis of chiral terminal (*E*)-1,3-enynes derived from the aldehydes (1*R*)-(−)-myrtenal and (*S*)-(−)-perillaldehyde using the alkynyl–phosphonio complex $[Ru\{C{=}CCH_2(PPh_3)\}(\eta^5\text{-}C_9H_7)(PPh_3)_2][PF_6]$ as synthon, 39
Epoxidation
Chiral bis(oxazoline) and pyridyl alcoholate dioxo-molybdenum(VI) complexes: synthesis, characterization and catalytic examinations, 207

1,1′-Ferrocenediyldialkoxysilanes
Synthesis of poly(ferrocenyldialkoxysilane) polymers and their introduction into oxide matrices, 46
Ferrocenyl diphosphines
Tunable ferrocenyl diphosphine ligands for the Ir-catalyzed enantioselective hydrogenation of *N*-aryl imines, 34
Fluorinated ligands
Highly efficient enantioselective catalysis in supercritical carbon dioxide using the perfluoroalkyl-substituted ligand (*R,S*)-3-H^2F^6-BINAPHOS, 130
Fluorinating reagents
Preparation of bistrimethylsilylmethylniobiumtetrafluoride and the application of KHF_2 and $n\text{-}Bu_4NHF_2$ as fluorinating reagents, 310
Functional phosphines
Chelating behaviour of a phosphinoalcohol leading to a stable alcohol palladium complex, 218

Gallium
Synthese und Charakterisierung chiraler Organogallium- und -aluminiumalkoholate mit intramolekularer Stickstoffstabilisierung. Röntgenstrukturanalyse von Dimethylgallium-(*S*)-2-dimethylamino-3-methyl-1-butanolat, Dimethylgallium-(*S*)-2-dimethylamino-3-phenyl-1-propanolat und Dimethylgallium-(*R*)-2-dimethylamino-1-butanolat, 317
Group 4 and 5 metals
Preparation of bistrimethylsilylmethylniobiumtetrafluoride and the application of KHF_2 and $n\text{-}Bu_4NHF_2$ as fluorinating reagents, 310

Hafnium
Preparation and crystal structures of the complexes $(\eta^5\text{-}C_5H_3TMS\text{–}CMe_2\text{–}\eta^5\text{-}C_{13}H_8)MCl_2$ and $[3,6\text{-}di^tButC_{13}H_6\text{–}SiMe_2\text{–}N^tBu]MCl_2$ (M = Hf, Zr or Ti): mechanistic aspects of the catalytic formation of a isotactic–syndiotactic stereoblock-type polypropylene, 267
Half-metallocene fragments
'*Supine*' or '*prone*' ligands: geometric preference of conjugated diene, 1-azadiene, and 1,4-diazadiene ligands on half-metallocene complexes of early transition metals, 224
High performance liquid chromatography
High performance liquid chromatography for facile analytical separation of the enantiomers of chiral organometallic complexes, 242
Homogeneous catalysis
Highly efficient enantioselective catalysis in supercritical carbon dioxide using the perfluoroalkyl-substituted ligand (*R,S*)-3-H^2F^6-BINAPHOS, 130
Homogeneous Ziegler–Natta catalysts
Preparation and structural features of 1-(2-pyridyl)- and 1-(2-furyl)indenyl-zirconium complexes, 197
Hydridoaluminates
A study of Noyori's reagent, 231
Hydrodesulfurization
Reactions of Cp*Ir(2,5-dimethylthiophene) with electrophiles, acids, and maleic anhydride, 55
Hydroformylation
Catalytic conversions in water: Part 19. Smooth hydroformylation of *N*-allylacetamide in mono- and biphasic aqueous media, 337

Hydrogen–deuterium exchange
Aqueous micellar and non-micellar effects during the asymmetric hydrogenation of dehydroamino acid derivatives: influence of amphiphiles on enantioselectivity and α-CH CD exchange, 158

Indenyl
Stereoselective synthesis of chiral terminal (*E*)-1,3-enynes derived from the aldehydes (1*R*)-(−)-myrtenal and (*S*)-(−)-perillaldehyde using the alkynyl–phosphonio complex [Ru{C≡CCH$_2$(PPh$_3$)}(η^5-C$_9$H$_7$)(PPh$_3$)$_2$][PF$_6$] as synthon, 39
Indenyl lithium compounds
Preparation and structural features of 1-(2-pyridyl)- and 1-(2-furyl)indenyl-zirconium complexes, 197
Insertion reactions
Insertion of carbon dioxide and isocyanide into tantalum–amide and tantalum–methyl bonds, 299
Intramolecularly donor-stabilized
Intramolecularly donor-stabilized silenes: Part 3. Synthesis and structure of 1-[2,6-bis(dimethylaminomethyl)phenyl]-1,2,2-tris(trimethylsilyl)silene, 261
Ir-diphosphine catalysts
Tunable ferrocenyl diphosphine ligands for the Ir-catalyzed enantioselective hydrogenation of *N*-aryl imines, 34
Iridium
High performance liquid chromatography for facile analytical separation of the enantiomers of chiral organometallic complexes, 242
Metallkomplexe mit biologisch wichtigen Liganden, CXXVII: Halbsandwich-Komplexe von Ruthenium, Rhodium und Iridium sowie phosphanhaltige Palladium- und Platin(II)-Komplexe mit Schiff-Basen aus α-Aminosäureanionen und Aldehyden, 109
Iridium complexes
Reactions of Cp*Ir(2,5-dimethylthiophene) with electrophiles, acids, and maleic anhydride, 55

Lanthanides
Cyclooctatetraene complexes of yttrium and the lanthanides with chiral and achiral aminotroponiminates, 277

Maleic anhydride
Reactions of Cp*Ir(2,5-dimethylthiophene) with electrophiles, acids, and maleic anhydride, 55
Mechanistic investigations
Aqueous micellar and non-micellar effects during the asymmetric hydrogenation of dehydroamino acid derivatives: influence of amphiphiles on enantioselectivity and α-CH CD exchange, 158
Metal complexes
Metallkomplexe mit biologisch wichtigen Liganden, CXXVII: Halbsandwich-Komplexe von Ruthenium, Rhodium und Iridium sowie phosphanhaltige Palladium- und Platin(II)-Komplexe mit Schiff-Basen aus α-Aminosäureanionen und Aldehyden, 109
Metallocene
Syntheses and properties of the novel co-catalysts *N,N*-dimethylanilinium- and trityl{tetrakis[4-(trifluoromethyl)-2,3,5,6-tetrafluorophenyl]borate}, 177
Micelles
Aqueous micellar and non-micellar effects during the asymmetric hydrogenation of dehydroamino acid derivatives: influence of amphiphiles on enantioselectivity and α-CH CD exchange, 158
Synthesis of achiral, but unsymmetric, seven-membered rhodium(I)-chelates for hydrogenation in the chiral environment of alkyl polyglucoside micelles, 120
Molybdenum
Chiral bis(oxazoline) and pyridyl alcoholate dioxo-molybdenum(VI) complexes: synthesis, characterization and catalytic examinations, 207
High performance liquid chromatography for facile analytical separation of the enantiomers of chiral organometallic complexes, 242
Mo$_2$(O^{i}Pr)$_4$(DMPE)$_2$
The electronic structure and bonding in W$_2$(μ-H)$_2$(O^{i}Pr)$_4$(DMPE)$_2$ and a comparison with Mo$_2$(O^{i}Pr)$_4$(DMPE)$_2$, 66

N-Allylacetamide
Catalytic conversions in water: Part 19. Smooth hydroformylation of *N*-allylacetamide in mono- and biphasic aqueous media, 337
N-Aryl imines
Tunable ferrocenyl diphosphine ligands for the Ir-catalyzed enantioselective hydrogenation of *N*-aryl imines, 34
^{n}Bu$_4$N$^+$ template
Self-assembly of [Cu(CN)$_4$]$^{3-}$ ions with cationic {Me$_3$Sn}$^+$ or {Me$_3$Sn(CH$_2$)$_3$SnMe$_3$}$^{2+}$ fragments in the presence of a ^{n}Bu$_4$N$^+$ template, 254
N-donor ligands
Ligands with cycloalkane backbones VIII. Steric effects in the ligand accelerated enantioselective alkylation of benzaldehyde, 153
Neodymium
Komplexkatalyse: LVII. Vereinfachte Synthese des Nd(π-C$_3$H$_5$)$_3$·C$_4$H$_8$O$_2$ nach der Grignard-Methode und Darstellung der neuen Allylneodym(III)-Komplexe [Nd(π-C$_3$Me$_5$)$_2$(π-C$_3$H$_5$)$_2$·C$_4$H$_8$O$_2$] und [Nd(π-C$_3$H$_5$)Cl(THF)$_3$]B(C$_6$H$_5$)$_4$·THF als Präkatalysatoren für die stereospezifische Butadienpolymerisation, 327
N ligands
Cyclooctatetraene complexes of yttrium and the lanthanides with chiral and achiral aminotroponiminates, 277
NMR
A study of Noyori's reagent, 231
Cationic derivatives of RhCl[P(η^2-C$_3$H$_5$)$_3$]. An intramolecular Diels–Alder rearrangement in the tripodal ligand tri(1-cyclohepta-2,4,6-trienyl)phosphane, P(C$_7$H$_7$)$_3$, 166
Non-hydrolytic sol–gel condensation
Synthesis of poly(ferrocenyldialkoxysilane) polymers and their introduction into oxide matrices, 46
Olefinic phosphane complexes
Cationic derivatives of RhCl[P(η^2-C$_3$H$_5$)$_3$]. An intramolecular Diels–Alder rearrangement in the tripodal ligand tri(1-cyclohepta-2,4,6-trienyl)phosphane, P(C$_7$H$_7$)$_3$, 166
Olefin polymerization
Syntheses and properties of the novel co-catalysts *N,N*-dimethylanilinium- and trityl{tetrakis[4-(trifluoromethyl)-2,3,5,6-tetrafluorophenyl]borate}, 177
Trimethylammonium{tetrakis(pentafluorophenoxy)borate}, a safe way to synthesize novel co-catalysts for olefin polymerization, 184
Optically active complexes
Stoichiometric cyclotrimerisation of chiral alkynes at a ruthenium centre: preparation of optically active (η^6-arene)(η^4-cycloocta-1,5-diene)ruthenium(0) complexes, 246
Optically active ligand
Synthese und Charakterisierung chiraler Organogallium- und -aluminiumalkoholate mit intramolekularer Stickstoffstabilisierung. Röntgenstrukturanalyse von Dimethyl-gallium-(*S*)-2-dimethylamino-3-methyl-1-butanolat, Dimethylgallium-(*S*)-2-dimethylamino-3-phenyl-1-propanolat und Dimethylgallium-(*R*)-2-dimethylamino-1-butanolat, 317

Organic synthesis
Alkinstabilisierte monomere Organo-Kupfer(I)-Verbindungen als Organyl-Transfer Reagenzien, 143
Organo-copper(I)
Alkinstabilisierte monomere Organo-Kupfer(I)-Verbindungen als Organyl-Transfer Reagenzien, 143
Organometallic fluorides
Preparation of bistrimethylsilylmethylniobiumtetrafluoride and the application of KHF_2 and n-Bu_4NHF_2 as fluorinating reagents, 310
Organotin(IV)
Self-assembly of $[Cu(CN)_4]^{3-}$ ions with cationic $\{Me_3Sn\}^+$ or $\{Me_2Sn(CH_2)_3SnMe_2\}^{2+}$ fragments in the presence of a $^nBu_4N^+$ template, 254
Osmium
Dihydroxylation of olefins using air as the terminal oxidant, 70
Oxazoline
Chiral bis(oxazoline) and pyridyl alcoholate dioxo-molybdenum(VI) complexes: synthesis, characterization and catalytic examinations, 207
Oxidation
Dihydroxylation of olefins using air as the terminal oxidant, 70
Oxidative addition
Diastereoselective oxidative addition of cyclic thiosulfinates to platinum(0) compounds: chiral platinum(II) complexes with sulfenato ligands Crystal structures of S(O)–CH_2–CH(OAc)–CH(OAc)–CH_2–S and dppePt-[S(O)–$(CH_2)_4$–S]: Part 16. Metal complexes of functionalized sulfur-containing ligands, 352
Oxide complexes
Chiral bis(oxazoline) and pyridyl alcoholate dioxo-molybdenum(VI) complexes: synthesis, characterization and catalytic examinations, 207
Oxo ligand
Catalytic hydrogenation of aromatics under biphasic conditions: isolation and structural characterisation of the cluster intermediate $[(\eta^6\text{-}C_6Me_6)_2(\eta^6\text{-}C_6H_6)Ru_4(\mu_3\text{-}H)_4(\mu_3\text{-}OH)(\mu_3\text{-}O)]^+$, 103

Palladium
Chelating behaviour of a phosphinoalcohol leading to a stable alcohol palladium complex, 218
Metallkomplexe mit biologisch wichtigen Liganden, CXXVII: Halbsandwich-Komplexe von Ruthenium, Rhodium und Iridium sowie phosphanhaltige Palladium- und Platin(II)-Komplexe mit Schiff-Basen aus α-Aminosäureanionen und Aldehyden, 109
Pentafluorophenol
Trimethylammonium[tetrakis(pentafluorophenoxy)borate], a safe way to synthesize novel co-catalysts for olefin polymerization, 184
Pentamethylcyclopentadienyl
Reactions of Cp*Ir(2,5-dimethylthiophene) with electrophiles, acids, and maleic anhydride, 55
Phenolato ligand
Mono(cyclopentadienyl)titanium complexes containing a sulfide-bridged bis(phenolato) ligand. Molecular structure of Ti{2,2′-S(OC_6H_2-4-Me-6-^tBu)$_2$}(η^5-C_5H_5)Cl, 3
Photoreactions with $M_2(CO)_{10}$ (M = Mn, Re)
Improved synthesis of the radical source bis[2,2,6,6-tetramethylpiperidyl-(1)]-disulfide and photoinduced reactions with $M_2(CO)_{10}$ (M = Mn, Re), 173
Planar chiral
High performance liquid chromatography for facile analytical separation of the enantiomers of chiral organometallic complexes, 242
Platinum
Diastereoselective oxidative addition of cyclic thiosulfinates to platinum(0) compounds: chiral platinum(II) complexes with sulfenato ligands Crystal structures of S(O)–CH_2–CH(OAc)–CH(OAc)–CH_2–S and dppePt-[S(O)–$(CH_2)_4$–S]: Part 16. Metal complexes of functionalized sulfur-containing ligands, 352
Metallkomplexe mit biologisch wichtigen Liganden, CXXVII: Halbsandwich-Komplexe von Ruthenium, Rhodium und Iridium sowie phosphanhaltige Palladium- und Platin(II)-Komplexe mit Schiff-Basen aus α-Aminosäureanionen und Aldehyden, 109
Synthesis and characterization of platinum(II)-terminated dendritic carbosilanes: X-ray crystal structure of the model species [PtCl(C_6H_3{CH_2NMe_2}-2-$SiMe_2$-5)(PPh_3)], 190
Poly(ferrocenylsilanes)
Synthesis of poly(ferrocenyldialkoxysilane) polymers and their introduction into oxide matrices, 46
Polymerization
Komplexkatalyse: LVII. Vereinfachte Synthese des Nd(π-C_3H_5)$_3$·$C_4H_8O_2$ nach der Grignard-Methode und Darstellung der neuen Allylneodym(III)-Komplexe [Nd(π-C_5Me_5)-(π-C_3H_5)$_2$·$C_4H_8O_2$] und [Nd(π-C_3H_5)Cl(THF)$_3$]B(C_6H_5)$_4$·THF als Präkatalysatoren für die stereospezifische Butadienpolymerisation, 327
Polymer properties
Unbridged metallocene dichloride complexes with mono-substituted indenyl ligands and their application for the polymerization of propene, 304
Polypropylene
Preparation and structural features of 1-(2-pyridyl)- and 1-(2-furyl)indenyl-zirconium complexes, 197
Potassium enolates
Meta, para and *meta, ortho* double *exo* nucleophilic additions of trimethylsilylester enolates derived from saturated and unsaturated carboxylic acids to tricarbonylchromium complexes of aryl ethers: dearomatizing cyclization to lactones, 284
Propene polymerization
Unbridged metallocene dichloride complexes with mono-substituted indenyl ligands and their application for the polymerization of propene, 304
2′-Pyridyl alcoholate
Chiral bis(oxazoline) and pyridyl alcoholate dioxo-molybdenum(VI) complexes: synthesis, characterization and catalytic examinations, 207
Rearrangements
Intramolecularly donor-stabilized silenes: Part 3. Synthesis and structure of 1-[2,6-bis(dimethylaminomethyl)phenyl]-1,2,2-tris(trimethylsilyl)silene, 261
Rhodium
Cationic derivatives of RhCl[P(η^2-C_7H_7)$_3$]. An intramolecular Diels–Alder rearrangement in the tripodal ligand tri(1-cyclohepta-2,4,6-trienyl)phosphane, P(C_7H_7)$_3$, 166
Metallkomplexe mit biologisch wichtigen Liganden, CXXVII: Halbsandwich-Komplexe von Ruthenium, Rhodium und Iridium sowie phosphanhaltige Palladium- und Platin(II)-Komplexe mit Schiff-Basen aus α-Aminosäureanionen und Aldehyden, 109
Rhodium(I) chelates
Synthesis of achiral, but unsymmetric, seven-membered rhodium(I)-chelates for hydrogenation in the chiral environment of alkyl polyglucoside micelles, 120
Rhodium/phosphine complexes
Catalytic conversions in water: Part 19. Smooth hydroformylation of *N*-allylacetamide in mono- and biphasic aqueous media, 337

Ruthenium
Catalytic hydrogenation of aromatics under biphasic conditions: isolation and structural characterisation of the cluster intermediate $[(\eta^6\text{-}C_6Me_6)_2(\eta^6\text{-}C_6H_6)Ru_3(\mu_3\text{-}H)_2(\mu_2\text{-}OH)(\mu_3\text{-}O)]^+$, 103
Metallkomplexe mit biologisch wichtigen Liganden, CXXVII: Halbsandwich-Komplexe von Ruthenium, Rhodium und Iridium sowie phosphanhaltige Palladium- und Platin(II)-Komplexe mit Schiff-Basen aus α-Aminosäureanionen und Aldehyden, 109
Stereoselective synthesis of chiral terminal (*E*)-1,3-enynes derived from the aldehydes (1*R*)-(−)-myrtenal and (*S*)-(−)-perillaldehyde using the alkynyl–phosphonio complex $[Ru\{C{=}CCH_2(PPh_3)\}(\eta^5\text{-}C_9H_7)(PPh_3)_2][PF_6]$ as synthon, 39
Stoichiometric cyclotrimerisation of chiral alkynes at a ruthenium centre: preparation of optically active (η^6-arene)(η^4-cycloocta-1,5-diene)ruthenium(0) complexes, 246
Schiff bases
Metallkomplexe mit biologisch wichtigen Liganden, CXXVII: Halbsandwich-Komplexe von Ruthenium, Rhodium und Iridium sowie phosphanhaltige Palladium- und Platin(II)-Komplexe mit Schiff-Basen aus α-Aminosäureanionen und Aldehyden, 109
Self assembly
Self-assembly of $[Cu(CN)_4]^{3-}$ ions with cationic $\{Me_3Sn\}^+$ or $\{Me_2Sn(CH_2)_3SnMe_2\}^{2+}$ fragments in the presence of a $^nBu_4N^+$ template, 254
Silaethenes
Intramolecularly donor-stabilized silenes: Part 3. Synthesis and structure of 1-[2,6-bis(dimethylaminomethyl)phenyl]-1,2,2-tris(trimethylsilyl)silene, 261
Silaheterocycles
Silaheterocyclen XXXVII. Die Kombination Alkenylhalogensilan/Lithium-*tert*-Butyl als Synthesebaustein zum Aufbau von Silaheterocyclen mit kumulierten CC-Doppelbindungen, 10
Silanes
Intramolecularly donor-stabilized silenes: Part 3. Synthesis and structure of 1-[2,6-bis(dimethylaminomethyl)phenyl]-1,2,2-tris(trimethylsilyl)silene, 261
Silenes
Intramolecularly donor-stabilized silenes: Part 3. Synthesis and structure of 1-[2,6-bis(dimethylaminomethyl)phenyl]-1,2,2-tris(trimethylsilyl)silene, 261
Silicon
Intramolecularly donor-stabilized silenes: Part 3. Synthesis and structure of 1-[2,6-bis(dimethylaminomethyl)phenyl]-1,2,2-tris(trimethylsilyl)silene, 261
Sol–gel process
Synthesis of poly(ferrocenyldialkoxysilane) polymers and their introduction into oxide matrices, 46
Stereospecific polymerization
Preparation and crystal structures of the complexes $(\eta^5\text{-}C_5H_3TMS\text{–}CMe_2\text{–}\eta^5\text{-}C_{13}H_8)MCl_2$ and $[3,6\text{-di}^tBuC_{13}H_6\text{–}SiMe_2\text{–}N^tBu]MCl_2$ (M = Hf, Zr or Ti): mechanistic aspects of the catalytic formation of a isotactic–syndiotactic stereoblock-type polypropylene, 267
Structure–activity correlation
Tunable ferrocenyl diphosphine ligands for the Ir-catalyzed enantioselective hydrogenation of *N*-aryl imines, 34
Substituted zirconocenes
Preparation and structural features of 1-(2-pyridyl)- and 1-(2-furyl)indenyl-zirconium complexes, 197
Substitution
Synthesis of chiral β-aminoalcohol-substituted carbene complexes of manganese and influence of the chiral carbene ligand on the diastereoselectivity of the CO/PR_3 exchange, 344
Sulfenato complexes
Diastereoselective oxidative addition of cyclic thiosulfinates to platinum(0) compounds: chiral platinum(II) complexes with sulfenato ligands. Crystal structures of $S(O)\text{–}CH_2\text{–}CH(OAc)\text{–}CH(OAc)\text{–}CH_2\text{–}S$ and dppePt-$[S(O)\text{–}(CH_2)_4\text{–}S]$: Part 16. Metal complexes of functionalized sulfur-containing ligands, 352
Sulfur donor
Mono(cyclopentadienyl)titanium complexes containing a sulfide-bridged bis(phenolato) ligand. Molecular structure of $Ti\{2,2'\text{-}S(OC_6H_2\text{-}4\text{-}Me\text{-}6\text{-}^tBu)_2\}(\eta^5\text{-}C_5H_5)Cl$, 3
Supercritical carbon dioxide
Highly efficient enantioselective catalysis in supercritical carbon dioxide using the perfluoroalkyl-substituted ligand (*R,S*)-3-H^2F^6-BINAPHOS, 130
Supine and *prone*
'*Supine*' or '*prone*' ligands: geometric preference of conjugated diene, 1-azadiene, and 1,4-diazadiene ligands on half-metallocene complexes of early transition metals, 224
Synthesis
Chelating behaviour of a phosphinoalcohol leading to a stable alcohol palladium complex, 218
Thiolato complexes
Diastereoselective oxidative addition of cyclic thiosulfinates to platinum(0) compounds: chiral platinum(II) complexes with sulfenato ligands. Crystal structures of $S(O)\text{–}CH_2\text{–}CH(OAc)\text{–}CH(OAc)\text{–}CH_2\text{–}S$ and dppePt-$[S(O)\text{–}(CH_2)_4\text{–}S]$: Part 16. Metal complexes of functionalized sulfur-containing ligands, 352
Thiophene
Reactions of Cp*Ir(2,5-dimethylthiophene) with electrophiles, acids, and maleic anhydride, 55
Thiosulfinates
Diastereoselective oxidative addition of cyclic thiosulfinates to platinum(0) compounds: chiral platinum(II) complexes with sulfenato ligands. Crystal structures of $S(O)\text{–}CH_2\text{–}CH(OAc)\text{–}CH(OAc)\text{–}CH_2\text{–}S$ and dppePt-$[S(O)\text{–}(CH_2)_4\text{–}S]$: Part 16. Metal complexes of functionalized sulfur-containing ligands, 352
Titanium complexes
Mono(cyclopentadienyl)titanium complexes containing a sulfide-bridged bis(phenolato) ligand. Molecular structure of $Ti\{2,2'\text{-}S(OC_6H_2\text{-}4\text{-}Me\text{-}6\text{-}^tBu)_2\}(\eta^5\text{-}C_5H_5)Cl$, 3
Transfer reagents
Alkinstabilisierte monomere Organo-Kupfer(I)-Verbindungen als Organyl-Transfer Reagenzien, 143
Tricarbonylchromium complexes
Meta, para and *meta, ortho* double *exo* nucleophilic additions of trimethylsilylester enolates derived from saturated and unsaturated carboxylic acids to tricarbonylchromium complexes of aryl ethers: dearomatizing cyclization to lactones, 284
Trimethylene tethers
Self-assembly of $[Cu(CN)_4]^{3-}$ ions with cationic $\{Me_3Sn\}^+$ or $\{Me_2Sn(CH_2)_3SnMe_2\}^{2+}$ fragments in the presence of a $^nBu_4N^+$ template, 254
Trinuclear
Catalytic hydrogenation of aromatics under biphasic conditions: isolation and structural characterisation of the cluster intermediate $[(\eta^6\text{-}C_6Me_6)_2(\eta^6\text{-}C_6H_6)Ru_3(\mu_3\text{-}H)_2(\mu_2\text{-}OH)(\mu_3\text{-}O)]^+$, 103

Tripodal ligands
Cationic derivatives of $RhCl[P(\eta^2\text{-}C_7H_{11})_3]$. An intramolecular Diels–Alder rearrangement in the tripodal ligand tri(1-cyclohepta-2,4,6-trienyl)phosphane, $P(C_7H_{11})_3$, 166

Unbridged metallocene complexes
Unbridged metallocene dichloride complexes with mono-substituted indenyl ligands and their application for the polymerization of propene, 304

Unsymmetric bisphosphines
Synthesis of achiral, but unsymmetric, seven-membered rhodium(I)-chelates for hydrogenation in the chiral environment of alkyl polyglucoside micelles, 120

Vinylidene complexes
Stereoselective synthesis of chiral terminal (*E*)-1,3-enynes derived from the aldehydes (1*R*)-(−)-myrtenal and (*S*)-(−)-perillaldehyde using the alkynyl–phosphonio complex $[Ru\{C{=}CCH_2(PPh_3)\}(\eta^5\text{-}C_9H_{11})(PPh_3)_2][PF_6]$ as synthon, 39

Water-soluble catalyst
Catalytic hydrogenation of aromatics under biphasic conditions: isolation and structural characterisation of the cluster intermediate $[(\eta^6\text{-}C_6Me_6)_2(\eta^6\text{-}C_6H_6)Ru_4(\mu_2\text{-}H)_2(\mu_2\text{-}OH)(\mu_3\text{-}O)]^+$, 103

Wittig reactions
Stereoselective synthesis of chiral terminal (*E*)-1,3-enynes derived from the aldehydes (1*R*)-(−)-myrtenal and (*S*)-(−)-perillaldehyde using the alkynyl–phosphonio complex $[Ru\{C{=}CCH_2(PPh_3)\}(\eta^5\text{-}C_9H_{11})(PPh_3)_2][PF_6]$ as synthon, 39

X-ray structure
Synthese und Charakterisierung chiraler Organogallium- und -aluminiumalkoholate mit intramolekularer Stickstoffstabilisierung. Röntgenstrukturanalyse von Dimethylgallium-(*S*)-2-dimethylamino-3-methyl-1-butanolat, Dimethylgallium-(*S*)-2-dimethylamino-3-phenyl-1-propanolat und Dimethylgallium-(*R*)-2-dimethylamino-1-butanolat, 317

X-ray structures
A study of Noyori's reagent, 231
Chelating behaviour of a phosphinoalcohol leading to a stable alcohol palladium complex, 218

Yttrium
Cyclooctatetraene complexes of yttrium and the lanthanides with chiral and achiral aminotroponiminates, 277

Zirconium
Preparation and crystal structures of the complexes $(\eta^5\text{-}C_5H_3TMS\text{–}CMe_2\text{–}\eta^5\text{-}C_{13}H_8)MCl_2$ and $[3,6\text{-di}^tBuC_{13}H_6\text{–}SiMe_2\text{–}N^tBu]MCl_2$ (M = Hf, Zr or Ti): mechanistic aspects of the catalytic formation of a isotactic–syndiotactic stereoblock-type polypropylene, 267
Syntheses and properties of the novel co-catalysts *N,N*-dimethylanilinium- and trityl{tetrakis[4-(trifluoromethyl)-2,3,5,6-tetrafluorophenyl]borate}, 177